

Package ‘TraitStats’

October 12, 2022

Title Statistical Data Analysis for Randomized Block Design Experiments

Version 1.0.2

Date 2022-06-28

Maintainer Nitesh Shirur Devaraja <mail@niteshgpb.in>

Description Functions for analysis of bulk data generated from experiments in Randomized block design as per Panse and Sukhatme (1954) <<https://books.google.co.in/books?id=Efo9AAAYAAJ>>. Computes analysis of variance; Descriptive statistics parameter like Mean, Minimum, Maximum, CV, Standard error of mean, Standard Error of deviation, CD; Genetic parameter statistics Genotypic Coefficient of Variation, Phenotypic Coefficient of Variation, Heritability in broad sense, Genetic Advance and Genetic Advance per cent mean; Variance and Co-variance matrix of genotypic, phenotypic and environmental; Correlation of genotypic, phenotypic and environmental. Further includes directly publication ready tables.

URL <https://www.niteshgpb.in/TraitStats.pdf>

License GPL-3

Encoding UTF-8

LazyData true

Imports utils, stats, lattice, rlist, grid, gridExtra, qpdf

Depends R (>= 3.5.0)

NeedsCompilation no

Author Nitesh Shirur Devaraja [aut, cre],
Parashuram Patroti [aut],
Shilpa Parashuram [aut],
Aravind J [ctb] (Reference material of augmentedRCBD),
Sunil Kumar [ctb]

Repository CRAN

Date/Publication 2022-06-30 16:10:02 UTC

R topics documented:

Barley	2
traitstats.envcorl	3
traitstats.envcov	4
traitstats.genocorl	5
traitstats.genocov	6
traitstats.genpar	7
traitstats.phenocorl	9
traitstats.phenocov	10
traitstats.rbd	11
traitstats.rbdmeantable	12
traitstats.rbdstats	13
traitstats.rbdsummary	14
Index	16

Barley	<i>Barley data (1973)</i>
--------	---------------------------

Description

Sample data on ear per plant, ear length (cms), 100-grain weight (g), and grain yield per plant (kg) from 1973 Barley performance trial, comprising of eight varieties in four replication (Chaudhary, 1973).

Usage

```
data("Barley")
```

Format

A data frame with 32 observations on the following 4 variables - EPR, EL,GW,and GY.

Source

Chaudhary BD. (1973). Estimation of genetic parameters in barley (*Hordeum vulgare* L.) by diallel analysis and its modifications. Unpublished Ph.D. Thesis, Haryana Agricultural University, Hisar.

Examples

```
data(Barley)
```

traitstats.envcor1 *Environmental Correlation Analysis*

Description

An extract of environment correlation coefficient for between traits

Usage

```
traitstats.envcor1(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

Environmental Correlation coefficient is calculated according to the Al-Jibouri et al., (1958).

Value

Environmental correlation coefficient matrix.

Author(s)

Nitesh, S.D., Parashuram Patroti, and Shilpa Parashuram.

References

Al-Jibour HA, Miller PA and Robinson HP. 1958. Genotypic and environmental variance in upland cotton cross of interspecific origin. *Agronomy Journal*, 50:633-637.

Examples

```
data("Barley")
traitstats.envcor1(Treatment=Barley$trt,
                  Replication=Barley$rep,
                  DataFile=Barley)
```

traitstats.envcov *Environmental Variance and Covariance*

Description

Computation of environment variance and covariance matrix.

Usage

```
traitstats.envcov(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

Environmental variance and covariance matrix is computed as per the methodology given by Al-Jibour et al., (1958).

Value

Environment variance and co-variance matrix. Diagonal value indicating the environmental variance and upper and lower half of the matrix indicating the environmental covariance.

Author(s)

Nitesh, S.D., Parashuram Patroti and Shilpa Parashuram

References

Al-Jibour HA, Miller PA and Robinson HP. 1958. Genotypic and environmental variance in upland cotton cross of interspecific origin. *Agronomy Journal*, 50:633-637.

Examples

```
data("Barley")
traitstats.envcov(Treatment=Barley$trt,
                  Replication=Barley$rep,
                  DataFile=Barley)
```

traitstats.genocor1 *Genotypic Correlation Analysis*

Description

An extract of genotypic correlation coefficient for between traits

Usage

```
traitstats.genocor1(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

Genotypic Correlation coefficient is calculated according to the Al-Jibouri et al., (1958).

Value

Genotypic correlation coefficient matrix.

Author(s)

Nitesh, S.D., Parashuram Patroti, and Shilpa Parashuram.

References

Al-Jibour HA, Miller PA and Robinson HP. 1958. Genotypic and environmental variance in upland cotton cross of interspecific origin. *Agronomy Journal*, 50:633-637.

Examples

```
data("Barley")
traitstats.genocor1(Treatment=Barley$trt,
                    Replication=Barley$rep,
                    DataFile=Barley)
```

traitstats.genocov *Genotypic Variance and Covariance*

Description

Computation of genotypic variance and covariance matrix.

Usage

```
traitstats.genocov(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

Genotypic variance and covariance matrix is computed as per the methodology given by Al-Jibour et al., (1958).

Value

Genotypic variance and co-variance matrix. Diagonal value indicating the genetic variance and upper and lower half of the matrix indicating the genotypic covariance.

Author(s)

Nitesh, S.D., Parashuram Patroti and Shilpa Parashuram

References

Al-Jibour HA, Miller PA and Robinson HP. 1958. Genotypic and environmental variance in upland cotton cross of interspecific origin. *Agronomy Journal*, 50:633-637.

Examples

```
data("Barley")
traitstats.genocov(Treatment=Barley$trt,
                  Replication=Barley$rep,
                  DataFile=Barley)
```

 traitstats.genpar *Genetic Parameter Analysis*

Description

traitstats.genpar performs genetic variability analysis for all the traits in the DataFile.

Usage

```
traitstats.genpar(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

traitstats.genpar performs genetic parameter analysis from the input values extracted from the ANOVA analysis for individual traits and computes several variability estimates. .

The phenotypic, genotypic and environmental variance (σ_p^2 , σ_g^2 and σ_e^2) are obtained from the ANOVA tables according to the expected value of mean square described by Federer and Searle (1976) as follows:

$$\sigma_p^2 = \text{Mean sum of squares of test treatments}$$

$$\sigma_e^2 = \text{Mean sum of squares of residuals}$$

$$\sigma_g^2 = \sigma_p^2 - \sigma_e^2$$

Phenotypic and genotypic coefficients of variation (*PCV* and *GCV*) are estimated according to Burton (1951, 1952) as follows:

$$PCV = \frac{\sigma_p^2}{\sqrt{\bar{x}}} \times 100$$

$$GCV = \frac{\sigma_g^2}{\sqrt{\bar{x}}} \times 100$$

Where \bar{x} is the mean.

The estimates of *PCV* and *GCV* are categorised according to Sivasubramanian and Madhavanmenon (1978) as follows:

CV (%)	Category
$x < 10$	Low
$10 \leq x < 20$	Medium
≥ 20	High

The broad-sense heritability (H^2) is calculated according to method of Lush (1940) as follows:

$$H^2 = \frac{\sigma_g^2}{\sigma_p^2}$$

The estimates of broad-sense heritability (H^2) are categorised according to Robinson (1966) as follows:

H^2	Category
$x < 30$	Low
$30 \leq x < 60$	Medium
≥ 60	High

Genetic advance (GA) is estimated and categorised according to Johnson et al., (1955) as follows:

$$GA = k \times \sigma_g \times \frac{H^2}{100}$$

Where the constant k is the standardized selection differential or selection intensity. The value of k at 5% proportion selected is 2.063. Values of k at other selected proportions are available in Appendix Table A of Falconer and Mackay (1996).

Value

Genotypic Coefficient of Variation and GCV Category
 Phenotypic Coefficient of Variation and PCV Category
 Heritability (broad sense) and h^2 Category
 Genetic Advance and GA Category
 Genetic Advance percent Mean and GAM Category

Note

Genetic parameter analysis need to be performed only if the sum of squares of treatment: Test is significant Negative estimates of variance components is computed are not abnormal. Ref. Dudley and Moll (1969).

Author(s)

Nitesh, S.D., Parashuram Patroti and Shilpa Parashuram

References

- Lush JH. (1940). Intra-sire correlations or regressions of offspring on dam as a method of estimating heritability of characteristics. Proceedings of the American Society of Animal Nutrition, 1940(1):293-301.
- Burtone GW and De Vane GM. (1953). Estimating heritability in tall Fescus (*Festuca arundinaceae*) from replicated clonal material. Agronomy Journal, 45:478-481.

Johnson HW, Robinson HF and Comstock RE. (1955). Estimates of genetic and environmental variability in soybeans. *Agronomy Journal*, 47:314-318.

Robinson HF. (1966). Quantitative genetics in relation to breeding on centennial of Mendelism. *Indian Journal of Genetics and Plant Breeding*, 171.

Dudley JW and Moll RH. (1969). Interpretation and Use of estimates of heritability of genetics variance in Plant Breeding. *Crop Science*, 9:257-262.

Sivasubramaniam S and Madhavamenon P. (1973). Genotypic and Phenotypic variability in rice. *The Madras Agricultural Journal*, 60:1093-1096.

Examples

```
data("Barley")
traitstats.genpar(Treatment=Barley$trt,
                 Replication=Barley$rep,
                 DataFile=Barley)
```

traitstats.phenocor1 *Phenotypic Correlation Analysis*

Description

An extract of phenotypic correlation coefficient for between traits

Usage

```
traitstats.phenocor1(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

Phenotypic Correlation coefficient is calculated according to the Al-Jibouri et al., (1958).

Value

Phenotypic correlation coefficient matrix.

Author(s)

Nitesh, S.D., Parashuram Patroti, and Shilpa Parashuram.

References

Al-Jibour HA, Miller PA and Robinson HP. 1958. Genotypic and environmental variance in upland cotton cross of interspecific origin. *Agronomy Journal*, 50:633-637.

Examples

```
data("Barley")
traitstats.phenocov1(Treatment=Barley$str,
                    Replication=Barley$rep,
                    DataFile=Barley)
```

traitstats.phenocov *Phenotypic Variance and Covariance*

Description

Computation of phenotypic variance and covariance matrix.

Usage

```
traitstats.phenocov(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

Phenotypic variance and covariance matrix is computed as per the methodology given by Al-Jibour et al., (1958).

Value

Phenotypic variance and co-variance matrix. Diagonal value indicating the phenotypic variance and upper and lower half of the matrix indicating the phenotypic covariance.

Author(s)

Nitesh, S.D., Parashuram Patroti and Shilpa Parashuram

References

Al-Jibour HA, Miller PA and Robinson HP. 1958. Genotypic and environmental variance in upland cotton cross of interspecific origin. *Agronomy Journal*, 50:633-637.

Examples

```
data("Barley")
traitstats.phenocov(Treatment=Barley$str,
                    Replication=Barley$rep,
                    DataFile=Barley)
```

traitstats.rbd	<i>RCBD Bulk Data Analysis</i>
----------------	--------------------------------

Description

Bulk RCBD data analysis for ANOVA and Descriptive statistics

Usage

```
traitstats.rbd(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

traitstats.rbd function analysis the input DataFile according to the methodology given by Panse and Sukhatme (1984). Futhermore, it also represent the analysis results along with descriptive statistics parameters like Trait mean, Minimum, Maximum, CV, StdError(m) and StdError(d) in the tabular representation of the publication ready format.

Value

Number of Replication
Number of Treatment
Treatment Mean Sum of Square along with Test of Significance
Replication Mean Sum of Square along with Test of Significance
Error Mean Sum of Square
Trait Mean
Minimum
Maximum
CV
StdError(m)
StdError(d)

Author(s)

Nitesh, S.D., Parashuram Patroti and Shilpa Parashuram

References

Panse and Sukhatme. 1984. Statistical Methods for Agricultural Workers. ICAR, New Delhi.

See Also

[traitstats.rbdmeantable](#), [traitstats.rbdstats](#), [traitstats.rbdsummary](#)

Examples

```
data("Barley")
traitstats.rbd (Barley$trt,
               Barley$rep,
               Barley)
```

```
traitstats.rbdmeantable
```

RCBD Mean Table

Description

Individual treatment mean table along with Grand trait mean, CV, SE(m) and CD.

Usage

```
traitstats.rbdmeantable(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

traitstats.rbdmeantable function analysis the input DataFile according to the methodology given by Panse and Sukhatme (1984) and generates the data frame consisting of the treatment mean over replication. Furthermore, it also represent the descriptive statistics parameter – Trait mean, StdError(m), CV and CD at 5 percent level of significance in the tabular representation of the publication ready format.

Value

Individual Treatment mean
Trait Mean
SE(m)
CV
CD

Author(s)

Nitesh, S.D, Parashuram Patroti and Shilpa Parashuram

References

Panse and Sukhatme. 1984. Statistical Methods for Agricultural Workers. ICAR, New Delhi.

See Also

[traitstats.rbd](#), [traitstats.rbdstats](#), [traitstats.rbdsummary](#),

Examples

```
data("Barley")
traitstats.rbdmeantable(Barley$str,
                        Barley$rep,
                        Barley)
```

traitstats.rbdstats *RCBD Descriptive Statistics*

Description

Descriptive Statistics for Trait Mean, Minimum, Maximum, SE(m), SE(d) and CV.

Usage

```
traitstats.rbdstats(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

traitstats.rbdstats function analysis the input DataFile according to the methodology given by Panse and Sukhatme (1984) and generates the RBD descriptive statistics for Trait Mean, Minimum, Maximum, Standard Error(mean), Standard Error(deviation) and Coefficient of Variation.

Value

Trait Mean
Minimum
Maximum
StdError(m)
StdError(d)
CV

Author(s)

Nitesh, S.D, Parashuram Patroti and Shilpa Parashuram

References

Panse and Sukhatme. 1984. Statistical Methods for Agricultural Workers. ICAR, New Delhi.

See Also

[traitstats.rbd](#), [traitstats.rbdmeantable](#), [traitstats.rbdsummary](#)

Examples

```
data("Barley")
traitstats.rbdstats(Barley$trt,
                   Barley$rep,
                   Barley)
```

traitstats.rbdsummary *RCBD Data Analysis Report*

Description

RCBD data analysis report in PDF formate compiling the ANOVA, Descriptive Statistics and Genetic Parameter along with Data distribution graph.

Usage

```
traitstats.rbdsummary(Treatment, Replication, DataFile)
```

Arguments

Treatment	Treatment column in the DataFile
Replication	Replication column in the DataFile
DataFile	Input Data file name

Details

`traitstats.rbdsummary` function analysis the input `DataFile` according to the methodology given by Panse and Sukhatme (1984) and generates the RCBD data analysis results comprising of Data point distribution, ANOVA table, Descriptive statistics and Genetic parameter arranged in the PDF format output file.

Value

Data distribution graph
ANOVA table : Rep, Trt, MStrt, MSrep and MSerror along the test of significance
Descriptive statistics table : Trait Mean, Min, Max, SE(m), SE(d), CV and CD.
Genetic Parameter : GCV, PCV, h², GA and GAM along with the category

Author(s)

Nitesh, S.D., Parashuram Patroti and Shilpa Prashuram

References

Panse and Sukhatme. 1984. Statistical Methods for Agricultural Workers. ICAR, New Delhi.

See Also

[traitstats.rbd](#), [traitstats.rbdmeantable](#), [traitstats.rbdstats](#)

Examples

```
data("Barley")
traitstats.rbd (Barley$trt,
               Barley$rep,
               Barley)
```

Index

Barley, [2](#)

traitstats.envcor1, [3](#)

traitstats.envcov, [4](#)

traitstats.genocor1, [5](#)

traitstats.genocov, [6](#)

traitstats.genpar, [7](#)

traitstats.phenocor1, [9](#)

traitstats.phenocov, [10](#)

traitstats.rbd, [11](#), [13–15](#)

traitstats.rbdmeantable, [12](#), [12](#), [14](#), [15](#)

traitstats.rbdstats, [12](#), [13](#), [13](#), [15](#)

traitstats.rbdsummary, [12–14](#), [14](#)